

- the treatments of varicose veins. *J Vasc Surg* 2011;53(5 Suppl):49S-65S.
2. Disselhoff BC, der Kinderen DJ, Kelder JC, Moll FL. Five-year results of a randomized clinical trial comparing endovenous laser ablation with cryostripping for great saphenous varicose veins. *Br J Surg* 2011;98:1107-11.
 3. Rasmussen LH, Bjoern L, Lawaetz M, Blemings A, Lawaetz B, Eklof B. Randomized trial comparing endovenous laser ablation of the great saphenous vein with high ligation and stripping in patients with varicose veins: short-term results. *J Vasc Surg* 2007;46:308-15.
 4. Rasmussen LH, Bjoern L, Lawaetz M, Lawaetz B, Blemings A, Eklof B. Randomised clinical trial comparing endovenous laser ablation with stripping of the great saphenous vein: clinical outcome and recurrence after 2 years. *Eur J Vasc Endovasc Surg* 2010;39:630-5.
 5. Rutherford RB, Padberg FT Jr, Comerota AJ, Kistner RL, Meissner MH, Moneta GL. Venous severity scoring: an adjunct to venous outcome assessment. *J Vasc Surg* 2000;31:1307-12.
 6. Garratt AM, Macdonald LM, Ruta DA, Russell IT, Buckingham JK, Krukowski ZH. Towards measurement of outcome for patients with varicose veins. *Qual Health Care* 1993;2:5-10.
 7. Smith JJ, Garratt AM, Guest M, Greenhalgh RM, Davies AH. Evaluating and improving health-related quality of life in patients with varicose veins. *J Vasc Surg* 1999;30:710-9.
 8. Perrin MR, Guex JJ, Ruckley CV, dePalma RG, Royle JP, Eklof B, et al. Recurrent varices after surgery (REVAS), a consensus document. REVAS group. *Cardiovasc Surg* 2000;8:233-45.
 9. Stonebridge PA, Chalmers N, Beggs I, Bradbury AW, Ruckley CV. Recurrent varicose veins: a varicographic analysis leading to a new practical classification. *Br J Surg* 1995;82:60-2.
 10. van den Bos R, Arends L, Kockaert M, Neumann M, Nijsten T. Endovenous therapies of lower extremity varicosities: a meta-analysis. *J Vasc Surg* 2009;49:230-9.
 11. Christenson JT, Gueddi S, Gemayel G, Bounameaux H. Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up. *J Vasc Surg* 2010;52:1234-41.
 12. Theivacumar NS, Dellagrammaticas D, Darwood RJ, Mavor AI, Gough MJ. Fate of the great saphenous vein following endovenous laser ablation: does re-canalisation mean recurrence? *Eur J Vasc Endovasc Surg* 2008;36:211-5.
 13. Winterborn RJ, Foy C, Earnshaw JJ. Causes of varicose vein recurrence: late results of a randomized controlled trial of stripping the long saphenous vein. *J Vasc Surg* 2004;40:634-9.
 14. Kostas T, Ioannou CV, Touloupakis E, Daskalaki E, Giannoukas AD, Tsetis D, et al. Recurrent varicose veins after surgery: a new appraisal of a common and complex problem in vascular surgery. *Eur J Vasc Endovasc Surg* 2004;27:275-82.
 15. van Rij AM, Jiang P, Solomon C, Christie RA, Hill GB. Recurrence after varicose vein surgery: a prospective long-term clinical study with duplex ultrasound scanning and air plethysmography. *J Vasc Surg* 2003;38:935-43.
 16. Sam RC, MacKenzie RK, Paisley AM, Ruckley CV, Bradbury AW. The effect of superficial venous surgery on generic health-related quality of life. *Eur J Vasc Endovasc Surg* 2004;28:253-6.
 17. Durkin MT, Turton EP, Wijesinghe LD, Scott DJ, Berridge DC. Long saphenous vein stripping and quality of life—a randomized trial. *Eur J Vasc Endovasc Surg* 2001;21:545-9.
 18. Michaels JA, Brazier JE, Campbell WB, MacIntyre JB, Palfreyman SJ, Ratcliffe J. Randomized clinical trial comparing surgery with conservative treatment for uncomplicated varicose veins. *Br J Surg* 2006;93:175-81.
 19. MacKenzie RK, Paisley A, Allan PL, Lee AJ, Ruckley CV, Bradbury AW. The effect of long saphenous vein stripping on quality of life. *J Vasc Surg* 2002;35:1197-203.

Submitted Oct 11, 2012; accepted Dec 13, 2012.

INVITED COMMENTARY

Joseph D. Raffetto, MD, *West Roxbury, Mass*

In this study, Rasmussen et al provide a 5-year follow-up of the original randomized trial in which endovenous ablation was compared with ligation and stripping of the great saphenous vein in symptomatic patients with varicose veins that was published after 2 years of follow-up.¹ The importance of long-term data demonstrating durability and patient satisfaction after venous surgery is imperative and was previously demonstrated by Merchant et al² in a multicenter prospective registry.

For many years, it has been presumed that endovenous ablation techniques should be the primary treatment modality for the refluxing great saphenous vein and completely replace ligation and stripping. This contention was speculative and without scientific evidence. A small randomized trial by Lurie et al³ in which endovenous radiofrequency ablation was compared with surgical ligation and stripping demonstrated that at 2 years of follow-up, in addition to similar treatment success and reduction in clinical severity, the endovenous ablation group had significantly reduced pain as a quality-of-life dimension compared with surgery that persisted up to 2 years.

Similar to the Lurie trial, the uniqueness of this study, however, is that it compares a well-validated surgical technique—ligation and stripping—with a relatively new treatment modality encompassing endovenous laser ablation, with important primary and secondary end points that mattered not only to the clinician but also to the patient and over a long period of post-treatment duration. Rasmussen et al found that both therapies were effective and had no differences in recurrent refluxing great saphenous vein, recurrent varicose veins, clinical severity, and quality-of-life analysis.

The strength in any study evaluating long-term results is the sample size in each group studied and the actual number of evaluable

patients at each year interval, and in this study, at the 5-year follow-up. In addition to patient recruitment, understandably not all patients will be available to evaluate, and in this study, the authors recognize these limitations. However, this should not take away from the significance of this well-designed study, with well-matched groups and relevant end points that are critically important in clinical practice for physician caring for patients with venous disease. Although our confidence in treating patients with varicose veins with less invasive modalities will increase, there is a barrage of other modalities that have not been studied in a carefully designed randomized trial that await both professional and clinical acceptance. I believe the study by Rasmussen et al has set the framework for further work in this important field, that will answer which treatments, which outcomes, and what follow-up is necessary in determining best treatment practices for patients with varicose veins.

REFERENCES

1. Rasmussen LH, Bjoern L, Lawaetz M, Blemings A, Lawaetz B, Eklof B. Randomized trial comparing endovenous laser ablation of the great saphenous vein with high ligation and stripping in patients with varicose veins: short-term results. *J Vasc Surg* 2007;46:308-15.
2. Merchant RF, Pichot O; Closure Study Group. Long-term outcomes of endovenous radiofrequency obliteration of saphenous reflux as a treatment for superficial venous insufficiency. *J Vasc Surg* 2005;42:502-9.
3. Lurie F, Creton D, Eklof B, Kabnick LS, Kistner RL, Pichot O, et al. Prospective randomised study of endovenous radiofrequency obliteration (closure) versus ligation and vein stripping (EVOLVEs): two-year follow-up. *Eur J Vasc Endovasc Surg* 2005;29:67-73.